



Topic	Pentacel®
Manufacturer	Sanofi Pasteur Detailed Prescribing Information (Fda.gov/media/74385/download)
Protects Against	Diphtheria, tetanus, pertussis, polio, and Haemophilus influenzae type b (Hib)
Routine Schedule	Four (4) dose series: 2, 4, 6, and 15-18 months
Minimum Intervals	4 week minimum interval between dose 1 and 2 4 week minimum interval between dose 2 and 3 6 month minimum interval between dose 3 and 4 (as determined by DTaP component)
Approved Ages	Children aged 6 weeks through 4 years
Administration	Intramuscular (IM) injection
Packaging	Vaccine is packaged as 5 single-dose vials 0.5mL of DTaP-IPV liquid component and 5 single-dose vials of lyopohilized Hib component
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) Do Not Freeze
Full ACIP Recommendations	Licensure of a Diphtheria and Tetanus Toxoids and Acellular Pertussis Adsorbed, Inactivated Poliovirus, and Haemophilus b Conjugate Vaccine and Guidance for Use in Infants and Children (ACIP) (Cdc.gov/mmwr/preview/mmwrhtml/mm5739a5.htm)
VFC Letter	VFC Clinical Letter (Eziz.org/PDF/VFC%20Providers-Pentacel%208-27-08%20v2%20FINAL.pdf)
Billing Codes	CHDP code: 82 CPT code for vaccine: 90698 CPT code for administration*: 90460, 90461 Medi-Cal Fee-For-Service (FFS) administration: 90698 with modifiers –SK (highrisk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23 Child Health and Disability Prevention (CHDP) Program Code Conversion (Files.medical.ca.gov/pubsdoco/chdp/articles/25768.02_Cd_Conv_Table.pdf)
Comments	Licensed in 2008