

Vaccine Fact Sheet: IPV



Topic	IPOL®
Manufacturer	Sanofi Pasteur Detailed Prescribing Information
Protects Against	Poliovirus
Approved Ages	Persons aged 6 weeks and older
CDPH Immunization Program Offering Product	 
Routine Schedule	Children: Four dose (4) series: 2, 4, and 6-18 months, and 4-6 years Adults: <u>Unvaccinated:</u> Three (3) dose series: 0, 1-2, and 6-12 months apart <u>Incompletely vaccinated (1-2 prior doses):</u> 1-2 doses to complete 3-dose series <u>Fully vaccinated (≥3 prior doses):</u> 1 lifetime booster if at increased risk of poliovirus exposure .
Minimum Intervals	Children: 4 week minimum interval between dose 1 and 2 4 week minimum interval between dose 2 and 3 6 month minimum interval between dose 3 and 4 Adults: 4 week minimum interval between dose 1 and 2 6 month minimum interval between dose 2 and 3
Administration	Intramuscular (IM) injection or Subcutaneous (SC) injection
Packaging	Vaccine is packaged as a 10-dose multi-dose vial. Use MDV polio until the expiration date once punctured.
Storage	Refrigerate between 36°F and 46°F (2°C to 8°C) Protect Vaccine from Light Do Not Freeze
Full ACIP Recommendations	Progress Toward Polio Eradication — Worldwide, January 2020–April 2022 MMWR (cdc.gov)
VFC Letter	Not available

Vaccine Fact Sheet: IPV (Continued)

Topic	IPOL®
Billing Codes	CPT code for vaccine: 90713 CPT code for administration*: 90460 Medi-Cal Fee-For-Service (FFS) administration: 90713 with modifiers –SK (high-risk) and –SL (VFC) ICD-10-CM code (encounter for immunization): Z23
Comments	Licensed in 1990

Refer to [Vaccine Fact Sheets](https://www.eziz.org/Resources/VaccineFactSheets) (EZIZ.org/Resources/VaccineFactSheets).