

Immunization Schedule with Combination Vaccines

	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	4-6 Years
Pediarix® Proquad® Quadracel™ or Kinrix®	Pediarix® DTaP, IPV, HepB + PCV Rotavirus Hib	Pediarix® DTaP, IPV, HepB¹ + PCV Rotavirus Hib	Pediarix® DTaP, IPV, HepB + PCV Rotavirus² Hib³	HepA MMR⁴ Varicella⁴ PCV⁵ Hib⁵	DTaP	HepA	Quadracel™ or Kinrix®⁸ DTaP, IPV + Proquad® MMRV
Pentacel®⁵ Proquad® Quadracel™ or Kinrix®	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus HepB	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus HepB¹	PENTACEL® DTaP, IPV, Hib + PCV Rotavirus² HepB	HepA MMR⁴ Varicella⁴ PCV⁵	PENTACEL® DTaP, IPV, Hib	HepA	Quadracel™ or Kinrix®⁸ DTaP, IPV + Proquad® MMRV
Vaxelis™⁷ Proquad® Quadracel™ or Kinrix®	Vaxelis™ DTaP, IPV, Hib, HepB + PCV Rotavirus	Vaxelis™ DTaP, IPV, Hib, HepB¹ + PCV Rotavirus	Vaxelis™ DTaP, IPV, Hib³, HepB + PCV Rotavirus²	HepA MMR⁴ Varicella⁴ PCV⁵ Hib⁵	DTaP	HepA	Quadracel™ or Kinrix®⁸ DTaP, IPV + Proquad® MMRV

Everyone 6 months+: 1) COVID-19 vaccine per current recommendations 2) Flu vaccine every fall⁶

1st RSV SEASON: RSV immunization for all infants < 8 months.⁹

Make sure the vaccine you administer contains the antigens on the doctor's order.

Keep it simple. Stick with the same product.

This is a suggested schedule for VFC providers ordering combination vaccines. For alternatives and details, consult the latest "Recommended Immunization Schedules for persons aged 0-18 years, United States." For more info, visit EZIZ.org.

1. A dose of Hepatitis B vaccine is not necessary at 4 months if doses are given at birth and 2 months but may be included as part of a combination vaccine.
2. The six month dose is not needed if Rotarix® was used exclusively for both dose 1 and 2 of the rotavirus vaccine series.
3. This six month Hib dose is not indicated if PedvaxHIB® is used exclusively for the 2 and 4 month infant doses.
4. CDC recommends MMR + Varicella at 12-15 months. Providers can use their discretion whether to use MMRV, however.
5. Can be administered as late as 15 months. For more information, consult the Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2023.
6. Influenza vaccine is available in thimerosal-free options. See California Health and Safety Code § 124172.
7. Licensed by FDA for children 6 weeks through 4 years of age (prior to the 5th birthday).
8. Licensed by FDA for children 4 through 6 years with previous doses of INFANRIX™ or PEDIARIX™. ACIP recommends that, whenever feasible, the same manufacturer's DTaP vaccines be used for each dose in the series; however, vaccination should not be deferred because the type of DTaP previously administered is unavailable or unknown. See CDC.gov/mmwr/preview/mmwrhtml/mm5739a4.htm.
9. Infants <8 months entering their first RSV season should receive RSV immunization if birth parent's prenatal RSV vaccination status is: unvaccinated, unknown, or vaccinated <14 days before birth. RSV immunization is also recommended for specified children ages 8-19 months who are at increased risk of severe RSV and entering their 2nd RSV season.



California
Vaccines for
Children Program